



C. Armitage Harper, III, M.D.
Jose Agustin Martinez, M.D.
James W. Dooner, M.D.
Mark Levitan, M.D.
Peter A. Nixon, M.D.
Robert W. Wong, M.D.
Shelley Day Ghafoori, M.D.
Ryan C. Young, M.D.

Patient Care Instructions Post Intravitreal Injection

Please follow these instructions after eye injections:

- Do not rub your eye after treatment
- For relief of mild to moderate pain or irritation, an over-the-counter pain reliever (i.e. Tylenol) or cool compress/ice pack can be used
- A new bottle of artificial tears/lubricating drops can be used
- No swimming or contact lens wear for 2 days
- It is okay to shower and wash your face. Keep eyes closed to avoid dirty water in the eyes.

The following symptoms are COMMON after successful eye injections for a day or so:

- Blurry vision
- Tearing
- Mild floaters or bubbles in your vision
- Mild to moderate discomfort or irritation
- Gritty, "Eyelash in the eye" sensation
- Redness (sometimes this can cover the entire white part of the eye and last a few weeks)
- Bloody tears for 1 - 2 days after treatment

The following symptoms are NOT NORMAL after eye injections, and you should call our office immediately if you experience:

- Severe, constant pain
- Worsening pain after the first day
- Significant decrease in vision
- Severe increase in floaters
- Curtain or veil in your vision

Please contact our office with any questions or concerns at (512) 451-0103 or 1-800-252-8259.

Follow-up care is very important. Be sure to keep your scheduled appointment.